

## MEDICATION DILUTION AND INJECTION INSTRUCTIONS

### **OVIDREL (hCG alpha) 250 mcg for ovulation**

1. Draw up 1cc of liquid from the diluent vial into your syringe
2. Place 1cc of diluent into powder vial and allow to dissolve completely
3. Draw the dissolved mixture into your syringe, replace the needle on the syringe with a subcutaneous needle (27ga, ½ inch)
4. Inject the medication subcutaneously as you were instructed

### **OVIDREL (hCG alpha) 125 mcg from an OVIDREL 250 mcg**

1. Draw up 1cc of liquid from the diluent vial into your syringe
2. Place 1cc of diluent into powder vial and allow to dissolve completely
3. Draw up ½ cc of the dissolved mixture into your syringe and place the remaining solution in the vial in the refrigerator for administration when directed
4. Replace the needle on your syringe with a subcutaneous needle ( 27ga, ½ inch) and inject the medication in your syringe as you were instructed

### **“COCKTAIL” MIXTURE USING OVIDREL 250mcg (you need 2 of these ) and GONAL-F, FOLLISTIM, OR REPRONEX (you need 2 of these)**

1. OVIDREL 375mcg from 2- OVIDREL 250mcg
  - a. Draw up 1cc of diluent into your syringe and place into one of the OVIDREL powders and allow it to dissolve completely, draw up ½ cc of mixture into your syringe and place remaining mixture in the vial in the refrigerator for administration when instructed
  - b. Take the mixture in your syringe and place in the second OVIDREL powder and allow it to dissolve, remove the empty syringe and go to second diluent vial and draw up ½ cc of diluent and place that in the mixture in the vial
  - c. Draw complete mixture back up into the syringe and replace the needle on the syringe with a subcutaneous needle (27ga, ½ inch) and inject the medication as you were instructed
2. GONAL-F, FOLLISTIM, OR REPRONEX (2 AMPS)
  - a. Draw up 1cc of diluent from one of the diluent vials into your syringe and place it first into one of the GONAL-F, FOLLISTIM, OR REPRONEX powders, allow it to dissolve completely and then draw it back into your syringe and place into the second powder and allow it to dissolve completely and draw the complete mixture into your syringe
  - b. Replace the needle on your syringe with a subcutaneous needle (27ga, ½ inch ) and administer the medication subcutaneously as you were instructed