

MEDICATION DILUTION AND INJECTION PROCEDURES

PROFASI (hCG) 10,000 units, PREGNYL (hCG) 10,000 units, NOVAREL(hCG) 10,000 units

1. Draw up 2cc of diluent from the diluent vial into your syringe
2. Place 2cc of diluent into vial of PROFASI, PREGNYL OR NOVAREL powder and allow it to dissolve completely
3. Draw all of the dissolved solution back into the syringe
4. Change the needle used for mixing to a new needle (21-22ga, 1 ½ inch) for intramuscular injection
5. Inject the medication intramuscularly as you were instructed

5,000 units of PROFASI, PREGNYL OR NOVAREL FROM A 10,000 unit of PROFASI, PREGNYL OR NOVAREL

1. Draw up 2cc of diluent from the diluent vial into your syringe
2. Place 2cc of diluent into the vial of PROFASI, PREGNYL, OR NOVAREL, 10,000 units and allow it to dissolve completely
3. Draw 1cc of the dissolved solution back into your syringe and leave the remaining 1cc in the vial and place that in the refrigerator to be used at the directed date
4. Replace the needle on the syringe with a new needle (21-22 ga, 1 1/2inch) for intramuscular injection
5. Inject the medication intramuscularly as you were instructed

MEDICATION MIXTURE FOR A “COCKTAIL” USING PROFASI, PREGNYL, OR NOVAREL 10,000 UNIT (2 are needed) AND GONAL-F, FOLLISTIM , HUMEGON, PERGONAL, OR REPRONEX(2 are needed of any one of these)

1. MIXTURE OF 15,000 UNITS OF PROFASI, PREGNYL OR NOVAREL
 - a. Draw up 2cc of diluent from liquid vial and place into one of the PROFASI, PREGNYL OR NOVAREL 10,000 powders, let the mixture dissolve and draw up 1cc into your syringe. Leave the remaining 1cc in the vial and refrigerate for use later at a directed date
 - b. Take the 1cc that is in the syringe and place into the second PROFASI, PREGNYL OR NOVAREL 10,000 powder, remove your syringe and go back to the diluent vial and draw up 1cc of liquid from the diluent vial and add that to the mixture in the vial, draw the complete mixture back up into your syringe
 - c. Change the needle on your syringe to a new needle (21-22ga, 1 ½ inch) for intramuscular injection as you were instructed
2. MIXTURE OF 2 AMPS OF GONAL-F, FOLLISTIM, HUMEGON, PERGONAL OR REPRONEX
 - a. Draw up 2cc of diluent from the diluent vial(s) into your syringe, place it first into one powder of GONAL-F, FOLLISTIM, HUMEGON, PERGONAL OR REPRONEX powder , let it dissolve and then draw it back into the syringe and place into the second powder and let it dissolve, draw it back up into the syringe
 - b. Change the needle on your syringe to a new needle (21-22ga, 1 ½ inch) for intramuscular injection as you were instructed